

# Vital fluorescent dyes for study of silicifying organisms

LIMNOLOGY  
FRESHWATER  
BIOLOGY

www.limnolfwbiol.com

Annenkov V.V.\*<sup>ORCID</sup>, Zelinskiy S.N.<sup>ORCID</sup>, Pal'shin V.A.<sup>ORCID</sup>, Avezova T.N., Danilovtseva E.N.<sup>ORCID</sup>

Limnological Institute, Siberian Branch of the Russian Academy of Sciences, Ulan-Batorskaya Str., 3, Irkutsk, 664033, Russia

**ABSTRACT.** The mini-review describes application of vital fluorescent dyes to study silica biosynthesis in diatom algae and siliceous sponges. Mechanisms of these processes are not clear and study of growing siliceous constructions in living organisms is complicated with silica transparency and presence of mature elements (siliceous valves of diatoms and spicules of sponges). Specific fluorescence dyes stain growing siliceous elements of skeletons and also primary siliceous particles which are formed during silicon capture from the environment. Vital fluorescent dyes based on 7-nitro-2,1,3-benzoxadiazole, pyridyl oxazole and coumarin fluorophores are described as well as examples of application of these dyes to study diatom algae and siliceous sponges.

**Keywords:** fluorescent dyes, diatom algae, siliceous sponges, biosilicification

## 1. Introduction

A lot of living organisms use siliceous constructions for building of important elements of their bodies. The most known are diatom algae (>20% of primary oxygen and organic carbon production (Treguer et al., 1995), siliceous sponges (water filterers and a source of biological active compounds), rise and horsetail. For many decades scientists try to answer question: how these organisms build highly ordered constructions from silica at ambient temperatures? This question is intriguing and important because:

- biosilica shows a great number of sophisticated constructions which are species specific (Fig. 1);
- biosilica properties are closer rather to amorphous quartz than to silica gel from aqueous medium (Grachev et al., 2008);
- living organisms synthesize siliceous constructions at ambient conditions, without high temperatures and aggressive chemicals;
- understanding of molecular mechanisms of biosilicification gives not only valuable information of these organisms but also provides new knowledge in silicon and organic chemistry including approaches to new materials.


Siliceous particles and constructions are colorless and transparent which complicates their study with optical methods. These objects are electron dense and

so are visible with transmittance electron microscopy but this method requires complex sampling and is not applicable for living specimens. Staining with vital fluorescent dyes is a powerful approach in biology which becomes more actual in last decades with dissemination of confocal microscopes. Rhodamines where the first dyes applied to staining growing cultures of diatoms (Li et al., 1989). These substances were accumulated in growing siliceous valves and girdle bands but the mature elements of siliceous frustules were colorless. The such behavior of rhodamines is explained with formation of silica in Silica Deposition Vesicles (SDVs) of diatoms. These intracellular organelles are close to lysosomes and their content is acidic (pH = 5.5) (Vrieling et al., 1999) which stimulate accumulation of amines in these vesicles. Application of rhodamines allowed to obtain interesting data on diatom physiology (Safonova et al., 2007) and to synthesize new fluorescent materials by biotechnological approach (Kucki and Fuhrmann-Lieker, 2012). Unfortunately, rhodamines are not the convenient instrument because of low gap between staining and toxic concentrations. On the other hand, quantum yield of rhodamines in aqueous medium is high (>70% (Arbeloa et al., 1988) which results in high fluorescence from the cultivation medium.

A set of new dyes for vital staining of silicifying organisms was elaborated and this mini-review describes compounds based on 7-nitro-2,1,3-benzoxadiazole (NBD) fluorophore and more specific dyes which change fluorescent spectrum after incorporation into silica (PDMPO and coumarin derivatives).

\*Corresponding author.

E-mail address: [annenkov@lin.irk.ru](mailto:annenkov@lin.irk.ru) (V.V. Annenkov)



**Fig. 1.** Scanning electron microscopy images of diatom siliceous valves (A – *Aulacoseira baikalensis* (K. Meyer) Simonsen 1979; B – *Stephanodiscus meyeri* Genkal & Popovskaya; C and D – *Ulnaria ferefusiformis* M. Kulikovskiy & H. Lange-Bertalot); E-G – sponge spicules of *Lubomirskia baicalensis* (Pallas, 1773). Scale bar represents 50 (E), 10 (A, C and F), 5 (G) and 2 (B, D)  $\mu\text{m}$ .


## 2. NBD based dyes for vital staining

The first NBD derivative for staining the growing siliceous frustules of diatoms was HCK-123 (Fig. 2) (Desclés et al., 2008). This dye contains a basic substituent which provides incorporation of the dye into acidic SDVs. NBD moiety is often applied for labeling of amine containing compounds (proteins, carbohydrates and etc.) with the use of 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole (Cl-NBD). This reagent is relatively cheap and readily reacts with primary and secondary amines.


Frustules of the diatom algae contain an interesting type of polyamines, so called long-chain polyamines (LCPAs). These compounds consist of amine groups separated by trimethylene fragments, with the first segment frequently containing four methylene groups (Kröger et al., 2000). Some amine groups are methylated, the LCPAs chains contain from several to

> 20 nitrogen atoms. These substances are available in very limited amounts from the natural sources which complicates study of their properties and role in biomineralization. We have elaborated (Annenkov et al., 2006; 2009) a stepwise approach to synthesis of LCPAs, including individual compounds with 2-7 nitrogen atoms and oligomeric mixture containing polyamines with > 20 nitrogen atoms.


The reaction of NBD-Cl with short polyamines (2-3 nitrogens) resulted in new dyes NBD-N2 and NBD-N3 (Fig. 2) (Annenkov et al., 2010). NBD-N3 as well as HCK-123 stains acidic organelles of eukaryotic cells but NBD-N2 is more selective and penetrates into growing siliceous structures only. This selectivity is explainable with shorter basic chain of NBD-N2 (one amine nitrogen only). NBD-N2 was successfully applied in study of initial stages of the diatom valve biosynthesis (Annenkov et al., 2013) and spiculogenesis



**Fig. 2.** Structures of NBD based dyes: HCK-123, NBD-N2 and NBD-N3.



**Fig. 3.** Initial stages of sponge spicule (A-F, *L. baicalensis*) and diatom valve (G and H, *U. ferefusiformis*) growth visualized by cultivation in the presence of NBD-N2 dye. The main stages of spiculogenesis in *L. baicalensis* are: silicon accumulation in sclerocyte (A, top); formation of organic filament and protrusion of the new spicule from the cell (A, bottom and B); further elongation of the filament, sclerocyte capture silicic acid and organic substances from the extracellular space which allows further growth of the spicule (C); new sclerocytes merge with the growing spicule (D and E) and a mature spicules (F) are obtained. Submicrometer siliceous particles are formed in the diatom cytoplasm after 30-60 s after silicon addition to silicon free cultural medium (G) and after 10-15 min new siliceous valve is visible (H). Scale bar represents: 20 (F) and 10 (A-E, G and H)  $\mu\text{m}$ .




**Fig. 4.** Fluorescent images of diatom siliceous valves after cultivation in the presence of NBD-N2 dye. A-D - natural samples obtained from Lake Baikal (Russia), E - siliceous valve of *U. ferefusiformis*. Diatoms were cultivated for 3 days in the presence of 0.5  $\mu\text{M}$  NBD-N2 and the biomass was treated with  $\text{H}_2\text{SO}_4/\text{H}_2\text{O}_2$  (1:1) mixture. Scale bar represents 10  $\mu\text{m}$ .


in sponge *L. baicalensis* (Annenkov and Danilovtseva, 2016). Fluorescent and confocal microscopy allowed to visualize primary submicrometer silicon-containing particles formed during capture of silicic acid with diatoms and growing spicules starting from single sponge cell (Fig. 3).

NBD-N2 and NBD-N3 penetrates into siliceous frustules of diatoms and sponge spicules (Fig. 4-6). The dyes are deeply buried in the biogenic silica and retain fluorescent activity after treatment with strong acids and oxidizing agents during cleaning of organic substances. The stained valves and spicules are suitable for study with confocal microscopy (Fig. 6) which allows to obtain additional information about internal structure of the material without real slicing.


**Fig. 5.** Fluorescent images of diatom siliceous valves and sponge spicules after cultivation in the presence of NBD-N2 dye. A - *L. baicalensis*, B - *Baikalospongia bacillifera* (Dybowsky, 1880), C - *Swartschewska papyracea* (Dybowsky, 1880) and D - *Baikalospongia intermedia* (Dybowsky, 1880). Sponges were cultivated for a month in the presence of 0.5  $\mu\text{M}$  NBD-N2 and the biomass was treated with bleach. Fluorescence allows to distinguish spicules grown during the experiment from old spicules. Scale bar represents 50  $\mu\text{m}$ .







**Fig. 8.** Siliceous materials stained with fluorescent dyes based on long chain polyamines. A - *L. baicalensis* spicules, B-D - Baikalean sediment. Scale bar represents 100 (A), 20 (B and C) and 10 (D)  $\mu\text{m}$ .





**Fig. 9.** Structure of Q-N2 and PDMPO dyes.

#### 4. Dyes with variable fluorescent properties

As discussed above, NBD based dyes stain acidic cell vesicles and SDV due to specific structure of the substituent at fluorophore group. The next generation of fluorescent dyes for vital staining of siliceous structures in living cells changes fluorescent spectrum under association with silica. The first representative of these dyes is 2-(4-pyridyl)-5-((4-(2-dimethylaminoethylaminocarbonyl)methoxy)-phenyl)oxazole (PDMPO, Fig. 9) (Shimizu et al., 2001; Parambath et al., 2016). Fluorescence of PDMPO depends on pH and the emission spectrum changes under association with silanol groups or involving into acidic vesicles. This behavior is useful in study of silicifying organisms as well as in study of silicification reactions *in vitro*.

Recently we have synthesized a new coumarin based fluorescent dye Q-N2 (Fig. 9) which contain amine group in the substituent (Annenkov et al., 2019). In contrast to PDMPO, fluorescence spectrum of this dye in aqueous medium does not depend on pH value. Aggregation of Q-N2 with silica or entrapping into siliceous materials results in enhancing of blue fluorescence and appearance of green emission. These effects were studied with fluorescence microscopy and they are easy visible with a routine epifluorescence microscope: fluorescence color changes from blue to cyan under excitation at 350-360 nm and green fluorescence appears under excitation at 470 nm (Fig. 10).



**Fig. 10.** Change of Q-N2 emission spectrum in the presence of 100 mM silicic acid during condensation at pH 7 (top spectrum - excitation at 365 nm, bottom spectrum - excitation at 470 nm), the data show faster growth of the fluorescence intensity in green area. The microscopy images show cells of *Aulacoseira* sp. after cultivation in the presence of 0.5  $\mu\text{M}$  Q-N2, top - light image, middle - excitation at 370 nm, bottom - excitation at 470 nm. Scale bar represents 10  $\mu\text{m}$ .


The absence of pH-depending changes of the Q-N2 fluorescence allows to attribute the strong cyan (green) fluorescence to siliceous structures. The other advantage of Q-N2 is the excitation maximum below 400 nm, far from excitation of chloroplasts. In combination with high quantum yield (11% comparing with <1% for NBD derivatives) this decreases damaging effect on living cells during microscopy investigation. Q-N2 shows inhibitory effect on the diatom growth in 10  $\mu\text{M}$  concentration (20-30% inhibition) which is 20 times higher its staining concentration. Q-N2 stains growing siliceous structures in diatoms and sponges (Fig. 11) and video monitoring of the silicon capture allowed us to find formation of condensed oligosilicates in several seconds after silicon capture with the diatom cell (Annenkov et al., 2019). This observation confirm hypothesis about silicon assimilation by diatoms in the form of partially condensed silicic acid.

## Acknowledgments

This work was supported by the Russian Academy of Sciences (project # AAAA-A16-116122110059-3); E. Danilovtseva thanks the Russian Foundation for Basic Research (Project # 16-04-00799 a). The authors wish to thank the Center of Ultramicroanalysis (Limnological Institute), the Baikal Analytical Center (Irkutsk Institute of Chemistry) and the Isotope-geochemical research center for Collective Use (A. P. Vinogradov Institute of Geochemistry of the Siberian Branch of the Russian Academy of Sciences) for providing equipment. We are deeply appreciative of I. Khanaev for the sponge samples and V. Itskovich for identification of the sponge species.

## References

- Annenkov V.V., Patwardhan S.V., Belton D. et al. 2006. A new stepwise synthesis of a family of propylamines derived from diatom silaffins and their activity in silicification. *Chemical Communications* 14: 1521–1523. DOI: 10.1039/b515967a
- Annenkov V.V., Zelinskiy S.N., Danilovtseva E.N. et al. 2009. Synthesis of biomimetic polyamines. *ARKIVOC* xiii: 116–130. DOI: 10.3998/ark.5550190.0010.d10
- Annenkov V.V., Danilovtseva E.N., Zelinskiy S.N. et al. 2010. Novel fluorescent dyes based on oligopropylamines for the in vivo staining of eukaryotic unicellular algae. *Analytical Biochemistry* 407: 44–51. DOI: 10.1016/j.ab.2010.07.032
- Annenkov V.V., Basharina T.N., Danilovtseva E.N. et al. 2013. Putative silicon transport vesicles in the cytoplasm of the diatom *Synedra acus* during surge uptake of silicon. *Protoplasma* 250: 1147–1155. DOI 10.1007/s00709-013-0495-x
- Annenkov V.V., Glyzina O.Yu., Verkhovzina O.N. et al. 2014. Fluorescent amines as a new tool for study of siliceous sponges. *Silicon* 6: 227–231. DOI 10.1007/s12633-014-9220-4
- Annenkov V.V., Verkhovzina O.N., Shishlyannikova T.A. et al. 2015. Application of 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole in analysis: fluorescent dyes and unexpected reaction with tertiary amines. *Analytical Biochemistry* 486: 5–13. DOI: 10.1016/j.ab.2015.06.025



**Fig. 11.** Initial stages of sponge spicules (A-E, *L. baicalensis*) and diatom valve (F and G, *U. ferefusiformis*) growth monitored with Q-N2 dye. The cyan color of new siliceous structures is distinguishable from blue fluorescence of free Q-N2. Scale bar represents 20 (A-E) and 10 (F and G)  $\mu\text{m}$ .

Annenkov V.V., Danilovtseva E.N. 2016. Spiculogenesis in the siliceous sponge *Lubomirskia Baicalensis* studied with fluorescent staining. *Journal of Structural Biology* 194: 29–37. DOI:10.1016/j.jsb.2016.01.010

Annenkov V., Danilovtseva E., Pal'shin V. et al. 2018. Fluorescently-tagged polyamines for the staining of siliceous materials. *Plant Physiology and Biochemistry* 125: 205–211. DOI: 10.1016/j.plaphy.2018.02.014

Annenkov V.V., Zelinskiy S.N., Pal'shin V.A. et al. 2019. Coumarin based fluorescent dye for monitoring of siliceous structures in living organisms. *Dyes and Pigments* 160: 336–343. DOI:10.1016/j.dyepig.2018.08.020

Arbeloa F.L., Ojeda P.R., Arbeloa I.L. 1988. Dimerization and trimerization of rhodamine 6G in aqueous solution. *Journal of the Chemical Society-Faraday Transactions II* 84: 1903–1912. DOI: 10.1039/F29888401903

Desclés J., Vartanian M., El Harrak A. et al. 2008. New tools for labeling silica in living diatoms. *New Phytologist* 177: 822–829. DOI: 10.1111/j.1469-8137.2007.02303.x

Grachev M.A., Annenkov V.V., Likhoshway Y.V. 2008. Silicon nanotechnologies of pigmented heterokonts. *BioEssays* 30: 328–337. DOI: 10.1002/bies.20731

Kucki M., Fuhrmann-Lieker T. 2012. Staining diatoms with rhodamine dyes: control of emission colour in photonic biocomposites. *Journal of the Royal Society Interface* 9: 727–733. DOI: 10.1098/rsif.2011.0424

Kröger N., Deutzmann R., Bergsdorf C. et al. 2000. Spe-

cies-specific polyamines from diatoms control silica morphology. *Proceedings of the National Academy of Sciences of the United States of America* 97: 14133–14138. DOI: 10.1073/pnas.260496497

Li C.-W., Chu S., Lee M. 1989. Characterizing the silica deposition vesicle of diatoms. *Protoplasma* 151: 158–163. DOI: 10.1007/BF01403453

Parambath M., Hanley Q.S., Martin-Martinez F.J. et al. 2016. The nature of the silicophilic fluorescence of PDMPO. *Physical Chemistry Chemical Physics* 18: 5938–5948. DOI: 10.1039/C5CP05105C

Treguer P., Nelson D.M., Van Bennekom A.J. et al. 1995. The silica balance in the world ocean: a reestimate. *Science* 268: 375–379. DOI: 10.1126/science.268.5209.375

Vrieling E.G., Gieskes W.W.C., Beelen T.P.M. 1999. Silica deposition in diatoms: control by the pH inside the silicon deposition vesicle. *Journal of Phycology* 35: 548–559. DOI: 10.1046/j.1529-8817.1999.3530548

Safonova T.A., Annenkov V.V., Chebykin E.P. et al. 2007. Aberration of morphogenesis of siliceous frustule elements of the diatom *Synedra acus* in the presence of germanic acid. *Biochemistry (Moscow)* 72: 1261–1270. DOI: 10.1134/S0006297907110132

Shimizu K., Del Amo Y., Brzezinski M.A. et al. 2001. A novel fluorescent silica tracer for biological silicification studies. *Chemistry & Biology* 8: 1051–1060. DOI: 10.1016/S1074-5521(01)00072-2